

# MONITORING OF EARTHQUAKES ACTIVITIES ALONG THE SYRIAN RIFT SYSTEM ( LEFT-LATERAL) BY USING REMOTE SENSING AND GIS DATABASE

MOUTAZ DALATI

General Organization for Remote Sensing ( GORS ), Damascus, Syria - dalati.m@mail.sy

**KEY WORDS:** Remote Sensing, Thematic, Geology, Data Structures, SAR, DEM/DTM, Interferometer.

## ABSTRACT:

Northwest of Syria is part of one of the very active deformation belt on the Earth today. This area and the western part of Syria are located along the great rift system (Left-Lateral or African- Syrian Rift System). Those areas are tectonically active and caused a lot of seismic events. The AL-Ghab Graben complex is situated within this wide area of Cenozoic deformation. The system formed, initially, as a result of the break up of the Arabian plate from the African plate since the mid-Cenozoic. The Arabian plate is moving in a NNW direction, whereas the African plate is moving to the North. The left-lateral motion along the Dead Sea Fault Zone accommodates the difference in movement rate between both plates. Some 80 km to the North of AL-Ghab Graben both plates move into the Anatolian plate, which causes the latter to move to the West. The analysis of TM Space Imagery and digital image processing of spectral data show that the lineaments along AL-Ghab Graben may be considered as linear Conjunctions accompanied with complex fracturing system. This complex is affected by distance stresses accompanied with intensive forces. This action indicates that these sites are active and in a continual movement. In addition to that, the statistic analysis of Thematic Mapper data and the features from a digital elevation model ( DEM ) produced from SAR interferometry show the existence of spectral structures at the same sites.

## INTRODUCTION




Figure 1, Showing the area under study ,the young fault-controlled relief of AL-Ghab Graben area.

The area under study is located in the Northwestern part of Syria (Figure 1). The AL-Ghab Graben is limited from the West by the steep slope of the Jabal An-Nusseiriyeh range and from the East by the similar steep but less high slope of the Jabal Az-Zawiyeh range (Figure 2). As a whole the Graben looks like a flat-bottomed ditch about 10 km wide. The Southern part of the Graben represents a flat marshy trough up to 15 km wide. The trough is filled with Pliocene lacustrine sediments. The Pliocene sediments are covered with a thin sheet of Quaternary lacustrine sediments. This part of the Graben is called the AL-Ghab trough. The occurrence of the Pliocene deposits in the trough is practically horizontal. The thickness of the Pliocene

lacustrine sediments was fully measured only at the Eastern slope in the Southern part of the trough. There it equals 90 m. In other places wells drilled down to depth of 150 m did not strike the basal part of the Tertiary lacustrine sediments. The Northern part of the Graben situated in the territory of Turkey opens into the depression of the Buheiret El-'Omqa lake. Here Pliocene marine deposits are developed. Unlike continental deposits, they are also spread in the uplifted block bordering the Graben in the West. In general, the beds in the Northern part of the Graben are gently dipping from the Eastern to Western flank of the structure. The AL-Ghab trough is separated from the Northern part of the Graben by the rise of the apex in the region of Kafar Dibbin. North of the trough, near the town of Jisr Ash-Shaghour, Pliocene lacustrine sediments appear from under Recent dried swamps and then Helvetian and Oligocene limestone crop out. The latter are exposed in the deep canyon of the Nahr El-Aasi River and compose the core of the Dibbin dome. The dome is somewhat elongated in a North-North-Western direction. The dip of the beds is gentle and usually does not exceed 5°. Only near the faults, which cut the structure at an oblique angle, inclinations reaching 30° are noted.

## REMOTE SENSING APPLICATIONS

The AL-Ghab Graben is reflected in the intensity of neotectonics affecting it. Neo-tectonics is active, such as active faults. TM optical and radar data are used to image, for example, fault scarps and actively growing folds. Visible and infrared imagery with moderate resolution is used to map lineaments, which are often surface expression of deep-seated fractures. The Interpretation of Space Imagery TM shows that the surface expression of AL-Ghab Basin is flat plain with almost no topographic relief. The plain is ~60 km long and ~15 km wide ( Figure 3 ). Faults would be expected along the western margin given a typical fault step-over arrangement for AL- Ghab Basin.




Figure 2, Showing the limitation of AL-Ghab Graben



Figure 3, TM image ( Colour composite ) covers AL-Ghab Graben . Date : 22 /08/1998 , Landsat -5 .

To the south of AL-Ghab graben , between two strands of the Dead Sea Fault System ( DSFS ) is the Missyf Graben (Figure.4). The eastern fault strand can be traced northward at the surface along the eastern margin of AL- Ghab Basin before bifurcating to the north-northeast.. The basic structure of AL-Ghab Basin is a fault-controlled double depocenter. The main depocenter is positioned beneath the southern portion of the

surface plain, and slight northward migration of that depocenter with time is clear from the seismic data( Figure 5). Also apparent are a mid-basin ridge and a second smaller depocenter to the north. Despite out interpretations that have used all available data, several issues regarding the evolution of AL-Ghab Basin remain unresolved. Furthermore, while the fault geometry controlling AL- Ghab Basin roughly fits the pattern of a 'step-over' basin), AL- Ghab Basin shows several departures from this simple transform-parallel extension case. Transverse structures, such as those found in AL- Ghab Basin, are also commonly observed in other strike-slip basins. The digital image processing of Radar imagery showing the presence of active and fresh faulting zones along the AL-Ghab Graben ( Figure 6 ). TM and SAR-DTM data, also showed a gradual color tone and interruptions of linear-ellipse shapes which reflecting the presence of discontinuity contours along the fault zone extension ( Figure 7 )




Figure 4 , Shows the AL-Ghab and Missyf Grabens and the Dead Sea Fault System ( DSFS )

This features refer to abundance of surface morphological features indicate to Fresh Faults. Recent faulting is expressed as freshly exposed soil within the colluvial apron visible by its light tone color. These indicators had been proved by field checks. Furthermore, the statistic digital analysis of the spectral data show that there are distribution of spectral plumes.

These plumes are decreasing in intensity and color contrast from the center of the site to the direction of its edges.




Figure 5, Migration of depocenter with time is clear from the seismic data

This proves that there are an active tectonic reflecting the behavior of the movement and earth stresses at these fracturing zones. Results from statistic analysis of DTM image( Figure 8 ), demonstrate that the restraining bend of AL-Ghab graben involves active strike slip faulting. The AL-Ghab graben appears capable of generating large earthquakes and it should be an essential element in any regional earthquake hazards Assessment ( Figure 9 ) .




Figure 8, AL-Ghab graben involves active strike slip faulting

AL- Ghab Basin is not a textbook example of a step-over basin. However, through comparison with other basin studies and models it was found that many of the second-order structures within AL-Ghab Basin are common to other strike-slip basins.

The basin asymmetry seen in AL-Ghab is probably related to the amount, and sense, of relative movement across the bounding lateral faults.




Figure 6, Shows the Fresh and Active Faults on Radar Image




Figure 7, Discontinuity contours along the fault zone extension / TM and SAR - DTM

### CONCLUTIONS

The results are consistent with the observed surface faults that show a greater amount of relative motion on the eastern basin-bounding strike-slip fault. Observations from AL- Ghab are echoed in theoretical models that show cross-basin oblique-slip faults accommodating initial basin opening, but most subsidence on the basin bounding faults. A northward shifting depocenter, and the subsequent development of a second depocenter in AL- Ghab Basin, are due to increasing fresh faults overlap with time and step-over of the lateral motion from the eastern to the western faults.

The system formed, initially, as a result of the break up of the Arabian plate from the African plate since the mid-Cenozoic (Figure 10 ). The tectonic evolution of Syria has been profoundly affected by movement on nearby Arabian plate boundaries, which almost completely surround the country. This deformation is evidenced in large-scale geologic structures within Syria, which include the Palmyrides folds(Figure 10). The northern continuation of the Dead Sea Fault into Lebanon and Syria is comprised primarily of the Serghaya ( SF ) and the Yammouneh ( YF ) faults that trace parallel and close lines in between the Lebanon and Anti-Lebanon mountain chains ( Figure 11 ). The SF is a branch of the main Dead Sea fault in southwestern part of Syria, while the YF merges with the AL-Ghab fault (GF) near the Bouqeah plain in western Syria. The south-north trending GF in turn extends parallel to the Syrian coast, and merges with the major East Anatolian Fault ( EAF ) in southern Turkey ( Figure 12 ). This complex system in Lebanon and Syria has been the location of numerous large historical earthquakes (  $M \geq 6.5$  ). Ambreasy and Barazangi ( 1989 ) reported that the fault zone may still be active and capable of generating more destructive earthquakes in the future with an estimated long return period of 200-350 years.




Figure 9




Figure 11




Figure 12




Figure 10

REFERENCES

Al-Maleh, K., 1976. Etude stratigraphique, petrographique, sedimentologique et geochimique du Cretace du N.W. Syrien (Kurd Dagh se environs d'Aafrine): Les aspects petroliers de la region. Universite Pierre et Marie Curie (Paris VI), Paris. 620 p.

BEICIP 1975. Gravity maps of Syria: Damascus (Syria). Bureau d'etudes industrielles et de cooperation de l'institut français du petrole, Hauts de Seine, 96 p.

Best, J.A., M. Barazangi, D. Al-Saad, T. Sawaf and A. Gebran 1993. Continental margin evolution of the northern Arabian platform in Syria. American Association of Petroleum Geologists Bulletin, 77, 173-193.

Beydoun, Z.R. 1999. Evolution and development of the Levant (Dead Sea Rift) Transform System: a historical-chronological review of a structural controversy. In C. Mac Niocaill and P.D. Ryan (Eds.), Continental Tectonics, Geological Society of London, special publications, 164, 239-255.

- Brew, G., R. Litak, M. Barazangi and T. Sawaf 1999. Tectonic evolution of Northeast Syria: Regional Implications and Hydrocarbon Prospects. *GeoArabia*, 4, 289-318.
- Chaimov, T., M. Barazangi, D. Al-Saad, T. Sawaf and A. Gebran 1990. Crustal shortening in the Palmyride fold belt, Syria, and implications for movement along the Dead Sea fault system. *Tectonics*, 9, 1369-1386.
- Chaimov, T., M. Barazangi, D. Al-Saad, T. Sawaf and A. Gebran 1992. Mesozoic and Cenozoic deformation inferred from seismic stratigraphy in the south-western intercontinental Palmyride fold-thrust belt, Syria. *Geological Society of America Bulletin*, 104, 704-715.
- Dalati, M. 2000 "lineaments on Landsat Images-Detection, and Tectonic significance in North Western depressions of Syria". Proceedings of XIX International Congress for photogrammetry and Remote Sensing, ISPRS, Volume XXXIII, Part B7/1, PP 301-308, Amsterdam The Netherlands
- Dalati, M. 1995 : " Application of Remote Sensing to Geology, Tectonic and Mineral Exploration, Case Study : El - Rouge depression, North west of Syria " proceedings of the Fourteenth International conference on Applied Geologic remote sensing, ERIM volume I, PP 411 - 419, Las Vegas, Nevada, U.S.A.
- Dalati, M. 1994 "Applications of Remote Sensing for Tectonic Purposes in El-Rouge Depression, North West of The Syrian Arab Republic ". Proceedings of the International Symposium on Spectral Sensing Research, ISSSR, Volume II, PP.859-867, San Diego, California, USA
- Dubertret L. 1933 Les formes structurales de la Syrie et de la Palestine, Leur Origine. *C. R. Acad. Scie.*, Paris, France, 195, P. 66
- Devyatkin, E.V., A.E. Dodonov, E.V. Sharkov, V.S. Zykin, A.N. Simakova, K. Khatib and H. Nseir 1997. The El-Ghab Rift Depression in Syria: Its Structure, Stratigraphy, and History of Development. *Stratigraphy and Geological Correlation*, 5, 362-374.
- Domas, J. 1994. The Late Cenozoic of the Al Ghab Rift, NW Syria. *Sborník geologických věd Antropozoikum*, 21, 57-73.
- Kent, W.N. and R.G. Hickman 1997. Structural development of Jebel Abd Al Aziz, northeast Syria. *GeoArabia*, 2, 307-330.
- Krenkel, E. 1924. Der Syrische Bogen. *Zentralblatt fuer Mineralogie, Geologie und Palaeontologie*, 9, 274-281.
- Lupa, J., 1999. Structure of the Ghab Basin and surroundings, Northwest Syria, derived from gravity modeling. Unpublished Thesis, Cornell University, Ithaca, NY, 152 p.
- Matar, A. and G. Mascle 1993. Cinématique de la Faille du Levant au Nord de la Syrie: Analyse Microtectonique du Fosse d'Alghab. *Geodinamica Acta*, 6, 153-160.
- Mouty, M. 1997. Le Jurassique de la Chaîne côtière (Jibal As-Sahilyeh) de Syrie: essai de biozonation par les grands foraminifères. *Comptes Rendus de l'Académie des Sciences de Paris/Sciences de la terre et des planètes*, 325, 207-213.
- Mouty, M., M. Delaloye, D. Fontignie, O. Piskin and J.-J. Wagner 1992. The volcanic activity in Syria and Lebanon between Jurassic and Actual. *Schweizerische Mineralogische und Petrographische Mitteilungen*, 72, 91-105.
- Trifonov, V.G., V.M. Trubikhin, Z. Adzhamyanyan, S. Dzhallad, Y. El' Khair and K. Ayed 1991. Levant fault zone in northeast Syria. *Geotectonics*, 25, 145-154.

