

## Establishing Global and Regional Geodetic Reference Frames: The International GPS Service for Geodynamics

Ivan I. Mueller (The Ohio State University, Columbus),  
James F. Zumberge, Ruth. E. Neilan (Jet Propulsion Laboratory, California Institute of Technology), Gerhard Beutler,  
Werner Gurtner (University of Berne)

**KEYWORDS:** International Service, GPS, Reference Frames, Orbits

### ABSTRACT

The International GPS Service for Geodynamics is an international scientific service which began formal operation at the beginning of 1994, following several years of planning [1] and a 1<sup>1</sup>/<sub>2</sub>-year pilot service. The IGS collects, archives, and distributes GPS observation data sets, and uses these to generate high accuracy GPS satellite ephemerides, earth rotation parameters, coordinates and velocities of the IGS tracking stations, GPS satellite and tracking station clock information, and atmospheric information.

To accomplish its mission, the IGS consists of a network of GPS tracking stations, Data Centers, Analysis Centers, an Analysis Coordinator, a Central Bureau, and a Governing Board. The accuracies of IGS products are sufficient to support current scientific objectives, including the realization of the International Terrestrial Reference Frame (ITRF), monitoring of the earth's rotation and deformation of its liquid and solid components, ionospheric monitoring, and scientific satellite orbit determination. A Central Bureau Information System (CBIS) allows public access to IGS products, which include precise GPS ephemerides within 2-3 weeks of real time. IGS contributors and customers communicate through electronic mail, and exchanges are archived for future reference.

### INTRODUCTION

The primary objective of the International GPS Service for Geodynamics is to provide a service to support, through GPS data products, geodetic and geophysical activities. Cognizant of the immense growth in GPS applications the secondary objective of the IGS is to support a broad spectrum of operational activities performed by governmental or commercial organizations. The service also develops the necessary standards and specifications and encourages international adherence to its conventions.

A proof of concept for the International GPS Service for Geodynamics was conducted with a three-month campaign during June–September 1992 [2], and was continued through a pilot service until the formal establishment of the IGS in 1993 by the International Association of Geodesy (IAG). The routine IGS started on January 1, 1994. IGS operates in close cooperation with the International Earth Rotation Service (IERS). The IGS *Terms of Reference* [3] describes in broad terms the goals and organization of the IGS.

The IGS collects, archives, and distributes GPS observation data sets of sufficient accuracy to satisfy the objectives of a wide range of applications and experimentation. These data sets are used by the IGS to generate data products, including high accuracy GPS satellite ephemerides, earth rotation parameters,

coordinates and velocities of the IGS tracking stations, GPS satellite and tracking station clock information, and ionospheric information.

In particular the accuracies of these products are sufficient for the improvement and extension of the International Terrestrial Reference Frame (ITRF) maintained by the IERS, the monitoring of solid earth deformations, the monitoring of earth rotation and variations in the liquid earth (sea level, ice-sheets, etc.), for scientific satellite orbit determinations, and ionosphere monitoring.

To accomplish its mission, the IGS consists of a network of GPS tracking stations, several Data Centers, Analysis Centers, an Analysis Coordinator, a Central Bureau, and a Governing Board.

### OPERATION OF THE IGS

The IGS global network today comprises of about 100 stations with GPS tracking receivers. Within the next few years, expansion of the global network will address the paucity of sites near longitude 90°E, with planned coverage in Central Asia and islands in the Indian Ocean. Regional densification networks are also being tied to the ITRF system, especially in the Arctic, Australia, Europe, East Asia, Japan, South America and the U.S.

Each site has a high-precision dual-frequency P-code receiver which records measurements at 30-sec intervals. IGS Operational Data Centers (Table 1) are in direct contact with the tracking sites, and are responsible for fetching raw receiver data, formatting it in a common standard [4], and forwarding the data to Regional or Global Data Centers.

Reformatted global tracking data from several Operational Data Centers are collected at Regional Data Centers. A local archive of the data received is maintained, and the data are transmitted to the Global Data Centers. Regional Data Centers thus serve to reduce traffic on electronic networks.

**TABLE 1. IGS DATA CENTERS**

<b>Operational &amp; Regional</b>	
National Oceanic and Atmospheric Administration	US
Institut für Angewandte Geodäsie	Germany
Natural Resources, Canada	Canada
Norwegian Mapping Authority	Norway
Jet Propulsion Laboratory	US
.....	
<b>Global</b>	
NASA/Goddard Space Flight Center	US
Institut Géographique National	France
Scripps Institution of Oceanography	US

The Global Data Centers serve as interfaces to the Analysis Centers and the external user community. Their purpose is to receive/retrieve, archive, and provide on-line access to tracking data. In addition, Global Data Centers provide access to Analysis Center products.

There are currently seven IGS Analysis Centers (ACs) (Table 2) – three in the US, one in Canada, three in Europe – that routinely analyze some subset of the data from the IGS global network. ACs compute precise GPS ephemerides and earth orientation parameters. Daily results are posted periodically, typically once each week.

The IGS Analysis Coordinator – at Natural Resources Canada – is responsible for producing the official IGS orbits/clock corrections based on the combination of orbits from the separate ACs [5]. Because of consistency checks, the combined orbit is largely free of some systematic errors that occasionally remain in results from individual ACs. Agreement among ACs in GPS satellite ephemerides is generally at the level of 10 to 30 cm. It is estimated that the absolute accuracy of the IGS orbits is at

least one order of magnitude better than the broadcast ephemeris, even when anti-spoofing is in effect.

IGS Rapid Orbits/Clocks are available about ten days. IGS Final Orbits are available after about one month. The two differ from each other mainly in the orientation of their reference frame. Lately, due to the improvement of data quality and distribution, the differences have been negligibly small for all practical purposes. For this reason as of July 1, 1996 the IGS Final Orbit series is planned to be discontinued and the Rapid Orbits will become the Final. At the same time daily IGS (Preliminary) Orbits/Clocks of somewhat lower accuracy will be made available to the user community.

**TABLE 2. IGS ANALYSIS CENTERS**

Astronomical Institute–University of Berne	Switzerland
European Space Agency	Germany
GeoForschungsZentrum	Germany
Jet Propulsion Laboratory	US
National Oceanic and Atmospheric Administration	US
Natural Resources Canada	Canada
Scripps Institution of Oceanography	US

Estimates of earth orientation and station coordinates from the ACs are coordinated with the IERS [6]. Through the IGS, GPS-derived station locations are contributing more and more to the ITRF. As mentioned above, a major goal of the IGS is to include in the ITRF the coordinates of a number of sites that comprise dense regional GPS networks. The computations of coordinates for these regional stations will be handled by a number of Regional Network Associate Analysis Centers (RNAACs) currently being appointed in several countries around the world.

Table 3 summarizes the approximate accuracies of IGS orbits, clocks, earth orientation, and station coordinates ([6], [7], [8]).

#### **CENTRAL BUREAU INFORMATION SYSTEM**

The Central Bureau Information System (CBIS) provides public access to products of the IGS, and also provides a means of electronic messaging among IGS participants. The system runs automatically.

The CBIS is a client on the Internet's World Wide Web, allowing access with user-friendly Mosaic and Lynx interfaces. For example, on most Unix systems, the command

**TABLE 3. APPROXIMATE ACCURACIES OF IGS PRODUCTS**

GPS ephemerides	5 – 30 cm
GPS clocks	0.5 – 5 nsec
Earth orientation (pole)	0.2 – 0.7 mas
Earth orientation (length of day )	0.1 – 0.5 ms/day
Station coordinates (annual)	3 mm – 3 cm

lynx <http://igscb.jpl.nasa.gov/>

will connect the user to the CBIS. For those with X-window terminals, `xmosaic` for Unix systems, or the equivalent Macintosh or Windows application, allows a graphical interface to the CBIS.

At the top level of the CBIS one views the following:

The Central Bureau, funded by NASA at Caltech's **Jet Propulsion Laboratory**, supports an Information Service:

- \* **IGS Mail**: An archive of exchanges among IGS participants.
- \* **IGS Reports**: An archive of reports from IGS Data and Analysis Centers.
- \* **Regional Mail**: An archive of exchanges among those interested in specific regional GPS networks.

These collections, with other data, are also available through **anonymous FTP**.

Categories in bold can be selected for further exploration. By sending a message to `igscb@igscb.jpl.nasa.gov`, you can request that your Internet e-mail address be added to one or more of the mail and report distribution lists. The **anonymous FTP** category is explained in more detail in Table 4, where top-level files and directories are given.

### BENEFITS TO USERS

The IGS has been in formal operation for over two years, and detailed analysis strategies that exploit the qualities of its data and products are still being developed.

High quality Rinex tracking data from the global tracking network, GPS ephemerides that are more accurate than the broadcast orbits by at least an order of magnitude, and precisely-determined coordinates of dozens (and eventually hundreds or more) of sites distributed over the

entire globe as well as regionally, all provide benefits to users.

**TABLE 4. TOP-LEVEL OF CBIS ANONYMOUS FTP AREA**

files:	IGSCB.DIR NEWS.TXT README.TXT TREE.TXT
center	contains information on the various IGS Centers (Analysis, Data, etc.)
data	indicates where – at which Data Centers – Rinex data reside
general	miscellaneous information on GPS, IAG, and IGS, including the IGS Terms of Reference
mail	contains indices of mail messages, and the messages themselves
product	contains Analysis Center products: precise ephemerides, earth orientation, weekly summaries
software	contains source code for CBIS-exploring utilities
station	contains information on IGS stations

Investigators who use GPS in local regions can include data from one or more nearby IGS stations, fix the site coordinates from such stations to their ITRF values, and fix GPS satellite positions to their IGS-determined values. By doing so the investigator can reduce data from his own network with maximum accuracy and minimum computational burden. Furthermore, the results will be in a well-defined global reference frame.

For more information contact

IGS Central Bureau  
 Jet Propulsion Laboratory 238-540  
 4800 Oak Grove Drive  
 Pasadena, CA 91109 USA

voice: 818 354 8330  
 FAX: 818 393 6686  
 e-mail: `igscb@igscb.jpl.nasa.gov`

### ACKNOWLEDGMENTS

The authors would like to thank all participants of the IGS for their contributions and continued dedication. These cooperative efforts have enabled the IGS to realize a successful and productive international operation.

## REFERENCES

- [1] Mueller, I. I., G. Beutler, The International GPS Service for Geodynamics – Development and Current Structure, *Proceedings of the Sixth International Geodetic Symposium on Satellite Positioning*, The Ohio State University, Columbus, March 1992, pp. 823-835.
- [2] Beutler, Gerhard, 1993, The 1992 IGS Test Campaign, Epoch '92, and the IGS Pilot Service: An Overview, in *Proceedings of the 1993 IGS Workshop*, G. Beutler, E. Brockmann, editors, March 25-26, 1993, University of Bern, Switzerland, pp. 3-9.
- [3] Mueller, I. I., International GPS Service for Geodynamics: Terms of Reference, in *Proceedings of the 1993 IGS Workshop*, G. Beutler, E. Brockmann, editors, March 25-26, 1993, University of Bern, Switzerland, pp. 10-14.
- [4] Gurtner, Werner, RINEX–The Receiver-independent Exchange Format, in *GPS World*, July 1994, v. 5, no. 7.
- [5] Beutler, Gerhard, Jan Kouba, Tim Springer, 1993, Combining orbits of the IGS processing centres, in *Proceedings of the IGS Analysis Centre Workshop*, Jan Kouba, editor, October 12-14, 1993, Natural Resources Canada, Ottawa, pp. 19-55.
- [6] Feissel, M., and N. Essaïfi, editors, *1993 IERS Annual Report*, July, 1994; Central Bureau of IERS, Observatoire de Paris, France
- [7] Jan Kouba, editor, *Proceedings of the IGS Analysis Centre Workshop*, October 12-14, 1993, Natural Resources Canada, Ottawa.
- [8] Blewitt, Geoffrey, Advances in Global Positioning System Technology for Geodynamics Investigations: 1978–1992, in *Contributions of Space Geodesy to Geodynamics: Technology*, David E. Smith, Donald L. Turcotte, Editors, Geodynamics Series volume 25, American Geophysical Union, Washington, D.C.