

COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER

George Hanuschak, Chief, Geospatial Information Branch
Rick Mueller, Remote Sensing/GIS Analyst
National Agricultural Statistics Service, U.S. Department of Agriculture
3251 Old Lee Highway, Fairfax, VA 22030
ghanuschak@nass.usda.gov
rmueller@nass.usda.gov

ABSTRACT

In the last several years, the National Agricultural Statistics Service (NASS) has developed a Cropland Data Layer (CDL) in geographic information system (GIS) format (See Figure 1) for public use. It is being used by over 1000 GIS proficient users for watershed monitoring, agribusiness planning, prairie water pothole monitoring, crop rotation pattern analysis and animal habitat monitoring and value added analysis by private sector remote sensing/GIS industry. The CDL can be viewed and ordered on CD at <http://www.nass.usda.gov/research/Cropland/SARS1a.htm>. The benefits to GIS proficient data users seem to easily outweigh the program costs. State governments especially are saving considerable resources by partnering with NASS in the Cropland Data Layer Program. As the number of States continue to expand, the costs per State will continue to decline and the benefits will be expanding with more data users and uses.

THE OLD ECONOMICS (1970's-1980's)

The economics has changed rather substantially over the last three decades. The initial statistics and software research in the early 1970's was dominated by research staff costs and computer processing costs. The first full State project, Illinois in 1975, cost \$750,000 but still included substantial research and statistical and software development costs. The first applications costs were established for processing the State of Iowa (1978) in time for end of season crop area estimation. At that time, the total cost per State was estimated at \$300,000 and the data source was Landsat Multi-Spectral Scanner (MSS) at 60-80 meter resolution, with 4 spectral bands of data.

By the early 1980's, there were eight States being processed. The total cost per State was \$150,000. Major item costs were associated with remote sensing analyst staff (although spread out over eight States which increased efficiency substantially over one or two States with about the same staffing levels) and the cost of processing the data on mini-computers and on the ILLIAC and Cray supercomputers. Data costs were \$300 per Landsat scene and generally a small portion of the total project cost. When Landsat thematic mapper became available in 1982 at 30 meter resolution with seven spectral bands of data, the program faced a fork (applications continued or a research program and switch to Landsat TM) in the road and only had resources to go in one chosen direction. At first, NASS continued the eight State application using Landsat MSS data, with a small amount of resources devoted to research on Landsat TM. Even with the limited research program resources, it became apparent that the accuracy levels could be improved substantially by using Landsat TM but at a higher cost for the data and for the information extraction (more bands and better resolution). Thus in 1988 due to budget cuts and U.S. government plans to not have MSS data on future Landsat or other earth resource observing satellites, NASS abandoned the eight State application using Landsat MSS but increased the research program with Landsat TM.

A new applications program was initiated in 1991 in Arkansas and Mississippi. In 1992 Louisiana was added to the program. Budget cuts hit again and Louisiana and Mississippi were dropped from the program. In fact in 1995, the program was only intact for Craighead county in Arkansas. Needless to say is that the program had a very rocky road at that point in time. Thus, it was apparent that NASS had to partner and resource share with other Federal and State government Agencies in order to continue and expand the program again and add a public cropland data layer for distribution as well as the internal statistical products. In preparation for this, NASS analyst staff emphasized a more automated and more expert system like version of PEDITOR to ease the labor burden on analysts. When this was adequately accomplished, NASS was in position to seek out Federal, State and University partnerships.

COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER

Resource Leveraging Partnerships

Starting in 1997, a new and key data license partnership between NASS and USDA's Foreign Agricultural Service (FAS) and the Farm Services Agency (FSA) was established. Landsat 5 data could be used for projects that serviced these agencies programs. Thus, the Landsat data costs for a program expansion were reduced. In addition, with Landsat 7, data costs at EROS Data Center were set at \$600 for a system geo-referenced data set without licensing restrictions. USDA FAS has a USDA-wide license for Landsat 7 with Radarsat for \$405 per scene. Thus, the Landsat data costs were no longer a major constraint as when the cost for Landsat 4&5 peaked for government buyers at \$4,400 per Landsat scene in the early 1990s.

The highest cost historically was for expert remote sensing and statistical analysts to run the complex PEDITOR package for both remote sensing outputs and for statistical estimates with relative sampling error outputs as well. NASS had a small group of expert analysts in its' Research Division who were doing centralized analysis for several States. However, it was recognized that a decentralized analysis staff directly in NASS State Statistical offices would expand the Agency's analytical capabilities. In addition the analyst will have the advantage of more localized knowledge of the crops and cropping practices and other sources of data to evaluate the relative contribution of the Landsat to the NASS crop area estimation program at the State and county level.

A local analyst (with a stable position and longevity) and State office management will be in a better position to service other State government and university and farm organizations and agribusinesses by providing them a Cropland Data layer in geographic information system (GIS) format. For example, the analyst in Illinois did both Illinois and Indiana in a regional concept which is the most cost efficient. Small two to four State regions will likely be the most cost efficient model.

This data layer has already proven valuable to data users of NASS who often combine the NASS cropland data layer with other data layers. Some examples are for water quality assessments for watersheds, and location plans for a new agribusiness facility, such as a soybean crushing plant and for grain storage and transportation planning and for prairie water pothole monitoring in North Dakota.

Since 1998, NASS has added the States of Illinois, Indiana, Mississippi, New Mexico, and Iowa and continued to do North Dakota and Arkansas as well. Florida A&M University and NASS also entered into a cooperative program for minority student outreach in remote sensing and GIS by analyzing a portion of northern Florida. Establishing and maintaining effective and win-win partnerships for the Cropland Data Layer is challenging though. The analyst position needs to be strongly supported by all the partnering organizations and needs some longevity (5 years or so) to be effective. Some of the partnerships are stronger and more stable in terms of longevity than others. A companion paper by Craig (ASPRS 2001) entitled "A Resource Sharing Approach to Crop Identification and Estimation" goes into more detail about each partnership for the Cropland Data Layer program.

THE NEW ECONOMICS (LATE 1990's AND 2000)

NASS has entered into a new economics for crop area estimation and a Cropland Data Layer for public consumption through the new resource leveraging partnerships. The total cost per State is now \$75,000 and should drop further with the addition of more States. See the attached Figures 2 and 3 for the graphic representation of the cost reduction over time (inflation adjusted and non-inflation adjusted). Major reductions were realized in the cost of the Landsat data and even more dramatic in the cost of data processing which has gone from mini-computers and supercomputers to high end PC desktops now. All the processes, including full Landsat scene multi-temporal pixel categorization, are done on PC's. The non-inflation adjusted total project cost has been cut in half (See Figure 2). When adjusting for inflation (See Figure 3), the cost reduction is more than fivefold from the first large scale application in the early 1980's. The current program and the early 1980's are the only comparable periods to compare costs over. The reason is that the number of States that labor costs are being spread over are very similar in number and scope of effort. Dramatic cost reductions have been achieved in the cost of the Landsat data and for the data processing of the Landsat data to categorize it into most likely crop types (See Figure 4). Spreading out the labor costs over as many States as possible and having co-funding partners is the key to future cost reductions.

Perhaps more importantly, there are now more accurate and valuable output products, both internal to NASS and external to NASS data users. These include a State level estimate with relative sampling error reductions on the order of two to five compared to the ground sample data only. County level (small area) estimates with measurable sampling errors which have been recently improved with several alternative estimators. Third, a new publicly available Cropland Data Layer at the 30 meter pixel level categorized by crop type and formatted for standard geographic information systems input.

COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER

The use of the Cropland Data Layer by GIS proficient data users outside of NASS is a major new thrust. In North Dakota, the NASS State Statistical Office (SSO), the North Dakota State University and an agri-business firm used the Cropland Data Layer as one input into a key agri-business decision. The location of a wheat strain processing plant was being evaluated. By combining the Cropland Data Layer with a transportation network layer, a decision model was built by the University and used in the agri-business decision for the plant location. Another anticipated major use of the Cropland Data Layer is in watershed monitoring and in fact is one of the major reasons for the North Dakota State government interest in the program. The State of Illinois chose to add to the Cropland Data Layer by adding other ground data on land covers and converting it to a Land Cover Layer.

Satellite Data Sources

NASS staff have a very strong preference for the Landsat data. The nadir look, the wide area image swaths, and the spatial resolution and spectral bands are virtually ideal for the NASS Cropland Data Layer and crop estimation program. The only issue with Landsat is the temporal frequency. NASS staff desires eight day coverage as a minimum and would prefer even better temporal coverage. Backup to the Landsat for the NASS program is the Indian IRS satellite series, which have many Landsat-like features. SPOT has not proven to be a cost effective backup for the NASS program. NASS has used SPOT for area sampling frame construction, on occasion, where price reductions are available through State consortiums or now under a USDA-wide license or existing credits from USDA/FAS.

NASS gets questions about its' plans for using very high resolution satellite data. At this time there are no major plans to use very high resolution data, such as IKONOS and QUICKBIRD in the NASS program other than for occasional small quality control checks. NASS has farmer reported data from a stratified area frame sample to evaluate the Landsat data. The swath width of IKONOS is too small for wall-to-wall coverage needed for the Cropland Data Layer and the cost per square mile too high relative to Landsat. There is one small scale research project in NASS and FAS for citrus tree counting where very high resolution data will be used at the request of the Florida Department of Citrus.

BENEFITS

The public use Cropland Data Layer GIS files are definitely of value to the GIS proficient data users who order them. It is difficult to put an exact, or even approximate, dollar value on the use of the files but here are some very important uses of the Cropland Data Layer. Among the important uses are: watershed and water quality monitoring (State of North Dakota government and the EPA and USDA and North Dakota State U.), grain transportation and storage planning, crop rotation pattern analysis across years, use in university teaching of remote sensing and GIS, interaction of soils and crops over large areas, further enhancements by the private sector remote sensing/GIS industry for custom solutions for their clients, quality control for other Landsat categorizations, official State of Illinois Land Characterization Map, animal habitat monitoring, prairie water pothole monitoring in North Dakota, evaluation of where to build a wheat straw processing plant in North Dakota, identifying catfish ponds in Mississippi, agribusiness planning for seed, fertilizer, pesticide, fungicide and crop insurance, etc.

Several of these applications have saved the user or generated additional revenue for the user in significant dollar terms. The States of Illinois and North Dakota, in particular, have saved considerable resources by partnering with NASS on the Cropland Data Layer. The uses and number of users is also growing rapidly and is still in the development stage. Again, while it is difficult to quantify the benefits to society, the authors believe that the benefits to date easily outweigh the program costs. In addition, NASS gets additional statistical inputs for crop county acreage estimates as well in a real time fashion. The Cropland Data Layer is the lowest level of aggregation that NASS provides and thus gives a very good look at the spatial distribution of crops, which is of considerable interest in the crop sector of the U.S. economy.

FUTURE PERSPECTIVE (2001-2010)

NASS will continue to pursue partnerships, primarily with State governments, to expand the crop area estimation and Cropland Data Layer program to more States. The desires are for program expansion to the top 15-20 total cropland States in the United States. The Midwest, the Delta and the Great Plains are the likely areas for

COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER

expansion as resources allow for. As the number of States continue to expand, the costs per State will continue to decline and the benefits will be expanding with more data users and uses.

ACKNOWLEDGEMENTS


The authors would like to extend their thanks and appreciation to all the analysts in the States involved. Those are Patrick Willis in Illinois, Dath Mita in North Dakota, and Ken Akins in Mississippi, and analysts in Washington, D.C., Mike Craig, Jim Mergerson, Charles Day and PEDITOR programmer Martin Ozga and ground data programmer, Jeff Johnson and Section Head, Robert Hale. In addition, the State Statisticians and their staffs for extensive ground data preparations for the States in the program. Thanks also go to Martin Ozga for recent improvements in PEDITOR aimed at reducing analyst labor, including a new mosaicking algorithm. Thanks to several members of the Area Frame Section and State office staffs for assisting in analysis preparations or analysis.

The authors would also like to extend their appreciation to Carol House, Fred Vogel, Joe Reilly, Rich Allen and Ron Bosecker of NASS for their headquarters executive management support of this program and to Allen Vandergriff of USDA/FAS and Glenn Bethel of USDA/FSA.

REFERENCES

- Allen, J. D., and Hanuschak, G. (1988). The Remote Sensing Applications Program of the National Agricultural Statistics Service: 1980-1987. SRB Staff Report Number SRB-88-08 USDA, NASS, Washington, DC-USA
- Bellow, M. (1994b). Large Domain Satellite Based Estimators of Crop Planted Area. In: *1994 Proceedings of the Section on Survey Research Methods*, August, 1994, American Statistical Association, Toronto, Canada, pp.194-199.
- Craig, M. (2001). A Resource Sharing Approach to Crop Identification and Estimation. *2001 ASPRS/ACSM Annual Convention Technical Papers, available on the CD, Proceedings of the ASPRS 2001 Conference*, Bethesda, MD-USA.
- Day, C. (2001). A Compilation of PEDITOR Estimation Formulas. RDD Research Paper RDD-01-01, USDA, National Agricultural Statistics Service, Washington, DC-USA.
- Fishman, J. (2001). The Landsat 7 Applications of Metapath Software International (MCI), USGS/NASA Landsat Data Continuity Workshop, Reston VA -USA.
- Graham, M. (1993). State Level Crop Area Estimation Using Satellite Data in a Regression Estimator. In: *Survey Methods for Businesses, Farms, and Institutions*, ICES Part I, NASS Research Report No. SRB-93-10, USDA, Washington, DC-USA, pp. 7-11.
- Hanuschak, G., Allen, R., and Wigton, W. (1982). Integration of Landsat Data Into The Crop Estimation Program of USDA's Statistical Reporting Service. *Proceedings of the 1982 Machine Processing of Remotely Sensed Data Symposium*, Eighth International Symposium, LARS, Purdue University, West Lafayette, IN-USA, pp. 45-56.
- Hui, Annie, Liew, Soo Chin, Kwoh, Leong, Keogn, Lim, Hock, Extraction and Utilization of Geometrical and Contextual Information in very High Resolution IKONOS Satellite Imagery, Centre for Remote Imaging, Sensing and Processing, National University of Singapore Blk., Singapore 119260, 2000
- Kay, Simon, Leo, Oliver, Peedell, Steve, Computer-assisted recognition of Oliver trees in digital imagery, Space Applications Institute, JRC of European Commission, Ispra, Italy, 2000.
- Mueller, R. (2000). Categorized Mosaicked Imagery from the National Agricultural Statistics Service Crop Acreage Estimation Program. Available on the CD: *Proceedings of the ASPRS 2000 Conference*, ASPRS, Bethesda, MD-USA.
- Ozga, M., and Craig, M. (1995). PEDITOR - Statistical Image Analysis for Agriculture. Presentation at the Washington Statistical Society (WSS) Seminar, April, 1995, USDA, National Agricultural Statistics Service, Washington, DC-USA.
- Ozga, M. (2000). Batch Processing of Remote Sensing Jobs on the PC. Available on the CD: *Proceedings of the ASPRS 2000 Conference*, ASPRS, Bethesda, MD-USA
- Vogel, F., and Bange, G. (1999). Understanding USDA Crop Forecasts. In: NASS/WAOB Miscellaneous Publication No. 1554, U.S. Department of Agriculture, Washington DC-USA.

Figure 1: *Cropland Data Layer
McLean County, Illinois*



COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER

Figure 2: *Cost Per State (Unadjusted)*






Figure 3: *Cost Per State (Adjusted for Inflation)*



*1983 adjusted to year 2000 dollars

Figure 4: Major Item Costs (Adjusted for Inflation)



COST AND BENEFIT ANALYSIS OF A CROPLAND DATA LAYER