

Introduction

Establishing protected areas is a conscious attempt to support the last remainder of a biodiversity which kept its natural properties in the current instable developing process. Applying scientific principles to study the bioenvironmental potentials and evaluating the ecologic potentials determines the land use type and the natural potential (ecologic potential) (Makhdom 1999). These studies have been already conducted in a traditional way by using the ground data, however, this method was a high cost, time consuming method and it does not work for hard accessible areas. As the result, remote sensed images are used to study the natural resources, forests, ranges, wild life, erosion and other bioenvironmental factors (Chaderi 1996). Consequently, studying the ecologic potentials of western habitats of Mazandaran, involving forests, ranges, apparent animal types such as some kinds of Mammals, birds, reptilian, etc. seems very necessary.

Materials and Methods

Ecologic resources include biological and physical sources, each of them involving different parameters. In general, 3 methods of identifying ecological sources include: 1- inventory and sampling 2- automatic optical interpretation of aerial photographs remotely sensed images and topographic maps 3- Geographic information system (Makhdom, 1999). At first, a supervised digital classification method was applied via a maximum likelihood classifier. To do this, seven groups of training samples were selected as a model of spectral reflectance of different phenomenon in the study area (including water, Garden, planted crops, unplanted crops, forest, pasture, rocky lands and manufactured areas) based on the ground survey data.

The resulting map has a total accuracy of 57.4% and a kappa coefficient of 72.3%. Then, an optical classification method was applied. To do this classification, the raster map was transferred to a vector one. Then, the obtained map was laid on SPOT 5 panchromatic image. Then, it was interpreted, reviewed and edited using ground data in an Arc view 3.2 environment. The resulting map had a total accuracy of 65.15% and a kappa coefficient of 78.65%. This matter indicates the high capability of spot5 data using a hybrid classification method to provide maps of the mountainous and non accessible areas. Ecological resources include:

Physical sources

Due to lack of a meteorology station in the study area, we used Synoptic station of Marivan and Paveh near Kusalan. Rivers and Brooklets were extracted from topographic maps and remote sensed images. Then, they were digitized. The registered points were transferred to GIS in order to identify the existing springs. Ground samples and geographic characteristics of springs obtained by GPS system were used. In the next stage, hydrographic maps of the Kusalan area were also provided.

Land Shape maps

To assess the parameters forming the land shape (height classes, slope classes and five main geographic directions) and to prepare the necessary maps, four sheets of topographic maps at the scale of 1:50000 of year 1997 were used. Finally, the land shape maps were obtained by overlaying the 3 map layers of height, slope and geographic direction.

Biological Resources

Plants

The extinction degrees of the plants in the study area were identified after careful frequent ground checking at different phenological stages of plant growth by collecting vegetation samples.

Animals

The information about the animals in the study area was collected by the wildlife habitat experts. To prepare the wildlife scattering map, the animal species index was determined. Then, habitat location was registered by GPS device and spatial data was transferred to GIS environment. The vector map of wildlife habitat was obtained through the hybrid interpretation of remotely sensed images.

Land shape

Since the land shape is a qualitative factor, remotely sensed images and a topographic map of 1:50000 associated with the field data were used to identify the scenery of Kusalan area.

Socio-economical studies

Conventional boundary map was prepared via hybrid interpretation of SPOT5 images and ground surveys with local people.

Evaluating the Bioenvironmental Potentials

The following processes are used in all methods of evaluating the bioenvironmental potentials:

- 1- Ecologic resources identification
- 2- Socio-economic data collection
- 3- Ecologic data collection and analysis
- 4- Ecologic potential evaluation for each unit comparing the ecologic properties of each unit with the model
- 5- Zoning
- 6- Providing the land use potentials in terms of the global criteria

Results

The results of bioenvironmental resource identification (ecologic and socio-economic resources) are as follows:

Ecologic resources

Physical resources

Results of studying the physical resources show:

Climate

Combination of climatic conditions of Marivan and Paveh stations, and averaging the climatic parameters shows the cold wet climate of Kusalan area.


Hydrology

There are abundant steep slope brooklets and springs in Kusalan habitat which all of them ends to Sirvan River. The main springs of Kusalan include: Hooseian, Lar, Odele, Hevar, Hevar Heye Shour.

Land shape

According the obtained results, many mountaintops and elevations with crest or round tops in the cornice or precipices shapes exist in Kusalan. The walls of valleys often have a steep or moderate slope, ending to bottom-line of brooklets. Direction of brooklets is mainly observed in linear form having a low width and a V shape profile view. The studies on topographic factors demonstrated the mountainous high altitude topography of Kusalan. Based on table 1 and map5, height variation range is 1800m and the main height classes are 1200-1800m in 40% of the study area. 27% of the area is covered with 2000-2500m classes which are suitable for protection according to evaluation models. Table 1 shows the percentage and area of each slope class. Distribution of classes is heterogeneous and the permanent predominant slope class is 30-60%. The slope above 60% covers 1/3 of the study area. Based on the results of this study, the area for the classes is north orientation (28/8%), west orientation (25/6%), east orientation (25/1%), and south orientation (20/2%). Due to the

uniformity of area distribution of the main orientations, the summer and winter time is proper for wildlife, so, animals won't migrate to the neighboring habitats. Land shape was determined by resultant of 3 physiographic factors of slope, height, orientation. As the result, 509 small ecosystems are obtained from Kusalan big ecosystem.



Area (%)	Area (ha)	Height from sea level(m)	Height class
17.4	4474.98	800-1200	1
40.6	10457.55	1200-1800	2
14	3652.2	1800-2000	3
26.9	6927.93	2000-2500	4
1	251.73	Over 2500	5

Biological resources

Plants

224 plant species were identified using field data and through a process of plant collection and identification in Kusalan. These identified species include: 78% forbs, 13% grasses, 8% trees and bushes and 1 % brushes. Table4 demonstrates the natural resource preservation value in Kusalan area (figures1-4). 30 species among these 224 species are exposed to extinction threats.



Figure1: Oak forests



Figure2: image1) *Medicago radiate*, 2) *Rheum Ribes*, 3) Down Tullip



Figure4:images1: *Alkana sp.*,2: *prangos sp.*,3: *Avena Sativa*, 4: *Bromus tectorum*



Figure 3: images1: *Legoecia cuminoides* ,2 (*Trifolium pilulare* ,)3)*Callipeltis cucularia*) ,4 *Bromus tomentellus*) 5 *Poa bolbosa* .

Growth form			Values(number of species)							Biologic form	
trees	bushes	grasses	brushes	forages	medicinal	industrial	ornamentals	genetics	Ecologic		protective
8	10	206	76	81	44	14	31	20	7	17	Plants


Animals

According to the studies on mammals, there are 23 species of them in Kusalan habitat. The most outstanding of them are: *Canis lupus*, *canis aureus*, *vulpes*, *Ursus*, *Mustela nivalis*, *Meles*, *Vormela peregusna*, *Lutra*, *Felis catus*, *Lynx*, *Felidae*, all of them belongs to Carnivora family.

Five family and subfamily of Rodents observed in this habitat, including: *Sciuridae*, *Criceinae*, *Gerbilinae*, *Muridae*, *Hystrigidae*.

Evaluating the ecologic potentials

This area was prepared for evaluation via ecologic resource identification and data collection and analysis. The evaluation was in terms of the ecologic resources (land shape, soil, vegetation cover, wildlife scattering, and other physical resources) playing a role in ecologic models and it was conducted through a multi-factor method according to an ecologic model. Six types of land use determined for the study area involving preservation, mass recreation, alternative recreation, forestry, gardening and agriculture (map6).



Zoning

Zoning was based on the ecologic and socio-economic factors. The results obtained by combining and overlaying the maps of ecologic potentials and conventional boundary, reveal 12978 biological units in Kusalan area. Combining the uniform units, we obtained 6 main zones: 1- safe zone, 2- protected zone, 3- mass recreation, 4- alternative recreation, and 5- cultural and historical zone, 6- reconstruction zone.

Area of each zone and its ratio to the total area were measured. zone 1(safe zone) covering 29% of the study area is extended over the eastern and western ranges of Doab River. Some parts of this zone are observed in southern range with a slope over 60%. The vegetation cover has a low density in this zone. Rocks are the main coverage of the area. Some other factors such as some threatened species of goat(main habitat) and *Ursus* have also affected on the selection of this zone. Zone 2 (preservation zone) with an area of 40/7% surrounds zone 1. This zone involves some parts of forests in the north and northeast ranges. *Lynx* habitat can be observed in this zone. 13/7% of the study area pertains to alternative recreation (zone 3), mainly observed in south ranges of Sirvan Permanent River and north ranges of Shahoo Mountain. Zone 4 (mass recreation) is located at north ranges with an area of about 3/21%. This zone was chosen because of the asphalt road of Marivan-Sanandaj and also the gravel rural road of Bahramabad –Bandol. Historical town of Horamaan with special customs for holding Pirshaliar ceremony is selected as zone 5 (cultural and historical zone). The reconstructive zone is 13/2%. Totally, 70/7% of the study area

has the potentiality to be preserved. In continue, the final zoning was done with regard to their natural condition, wetland shape and their combination with the conventional boundary map. According to the results of zoning, 29% of the study area pertains to zone 1 and 40/7% is considered as zone 2. In total, 70% of the study area has the potentiality to be preserved. About 17% of the study area (zones 3&4) has recreative potentiality. 14% of zone 5 and the rest 13/2% are considered as zone 6 or the reconstructive area.

Conclusion

Results of this study indicate potentiality of the study area (Kusalan) to be a national park and receive the IUCN criteria. In this study, it is also verified that the biological potentials of an area or a natural habitat can be studied using data extracted from SPOT5 remotely sensed images (panchromatic 10 m images) via optical interpretation and supervised digital classification (hybrid method) and based on the global criteria (IUCN) to establish protected areas. It is also emphasized on using GIS to save and analyze the special data and also on providing proper outputs (maps, tables, etc.) as a significant and applicable method.